


RECOMMENDED PRACTICES FOR SMALL HOLDER PIGGERY ENTERPRISE



Supported By



Designed & Developed by



HOUSING



Allot 10-12 sq. ft. area per animal



The floor should be sloped (5 degree) backward for easy removal of excreta




Sty must be cross-ventilated



Feeding pan must be more than 2 feet wide & 1-2 feet in length





MANAGEMENT OF PREGNANT SOW



Proper record of mating, mating partner, date of farrowing, litter size, date of weaning, birth weight of piglet should be maintained



Heat should be detected timely & mating should be made at peak of heat



Pregnant sow should be kept alone & fed with high protein & mineral diet




Pregnant sow should be kept in farrowing pan with area more than 20 sq ft










Keep checking pregnant sow around time of farrowing & assist her in case of any difficulty





PIGLET MANAGEMENT

-  The piglets should be fed on colostrum at the earliest
-  The naval cord must be dipped in some antiseptic solution
-  Piglets should be fed on sow milk at least 4 times a day till one-week age
-  After one-week age, piglets should be kept on high protein, cooked diet
-  The piglets should be weaned at 21- 35 days
-  The piglets should be administered 2 Iron injections at day 4 & 14 of their life (1 ml I/M)





FEEDING MANAGEMENT




Feed should be properly cooked & should be fed on the same day



Mineral Mixture (15-20gm/Kg.) & Common Edible Salt (10gm/Kg) must be added to the feed



A balanced mixture of Kitchen Waste, Raw Grains, Roots, Green leaves, Fish Meal/ Oil cake should be prepared





VACCINATION & DEWORMING





Sl. No.	Age	Vaccine	Route	Dose
1	30 days	Swine Fever	I/M or S/C	1 ml
2	45 days	FMD	I/M or S/C	1 ml
3	60 days	Swine Fever	I/M or S/C	1 ml
4	75 days	FMD	I/M or S/C	1 ml


Deworming (Fenbendazole- Oral)		
Sl. No.	Age	Dose
1	42 days	1 Tab (150 mg)
2	63 days	1 Tab (150 mg)
3	5 Month	2 Tab (150 mg)
4	8 Month	3 Tab (150 mg)
5	11 Month	4 Tab (150 mg)



An illustration of a pigsty. On the left, a person's legs and feet are visible, wearing blue trousers, white socks, and orange boots. They are using a long-handled white scraper to clean a dark, oval-shaped puddle on a floor made of dark grey wooden planks. Several pink pigs are scattered around the sty. Two large, irregular brown stains are on the floor. The background shows vertical wooden posts.

SANITATION, HYGIENE & BIO-SECURITY

-  The floor should be maintained dry
-  The feeder & floor should be cleaned daily using KMnO_4 (Potassium Permanganate) solution
-  Visit of outside persons shall be strictly prohibited near the pig sty
-  Smearing the animal with Ash or washing with Tobacco water shall help in getting rid of External Parasites










Supported by



Design & Developed by



RECORD KEEPING

-  Daily record of the feed consumption should be maintained
-  Weekly growth monitoring
Age, Vaccination & Deworming records should be properly maintained
-  Disease records should be maintained with treatment given & health improvement
-  Sale & Purchase record should be maintained properly with the name of supplier & Buyer, date of purchase & sale
-  Pregnant sow record should be maintained with the date of farrowing, litter size, date of weaning, birth weight of piglet

